




Address Correction Requested



No May Breeding Survey, What does it mean for limits?

The U.S. Fish and Wildlife Service was unable to perform the Waterfowl Breeding Population and Habitat Survey (WBPHS or May Survey) this year. The May Survey is used to set bag limits and advise management and correct population models for the next year. Similarly, the Canadian Wildlife Service and several state and provincial agencies also cancelled their participation for 2020. Decisions to cancel the May Survey and other migratory bird monitoring this spring were based on the priority of protecting the health and safety of the American public and employees. The severe domestic and international travel restrictions also made many operations impossible.



Strata and transects of the Waterfowl Breeding Population and Habitat Survey (yellow -- traditional survey area, green -- eastern survey area).

The Service gave serious consideration to implementation of partial surveys but ultimately concluded that the development of analytical procedures to inform 2021-2022 hunting season recommendations using the most current and complete data was a more defensible and reliable strategy given ongoing and unpredictable COVID-19 restrictions. The Service will work with the Flyway Councils and Canadian Wildlife Service to develop alternative, temporary methods for formulating harvest recommendations in the absence of spring 2020 breeding population data, including developing adjustments to Adaptive Harvest Management strategies for the 2021-2022 duck seasons. Adjustments to goose, Sandhill Crane, American Woodcock, and dove harvest strategies will also be necessary and are being considered.

Cancellation of these surveys will impact population estimates and harvest management decisions for most ducks, geese, and webless gamebird species such as American Woodcock, and the mid-continent population of Sandhill Cranes. The Service, in consultation with the Flyway Councils, will use long-term data from spring/summer monitoring for these species to make regulatory harvest management decisions.

For the general duck seasons, the Service uses the long-term data and models to predict 2020 spring abundances of ducks and habitat conditions in lieu of the spring 2020 data, which was not collected. The results from these predictions will be combined with the existing harvest strategies to determine appropriate levels of harvest for the 2021-2022 season. This will ensure the sustainability of ducks and provide hunting opportunities for the American public.

Due to the amount of time that it takes to process the huge amount of data collected from the breeding surveys, the season limits are based off of data from the previous year (2020 bag limits are based on the spring surveys from 2019).

Bag limits for the 2020 Pacific Flyway duck season have already been released, but limits for 2021 will be affected by the cancellation of the may 2020 survey.



Photo credit Andrea Mott, USGS

Reminder!

Diversions Reporting for the California Water Board is due by July 1st 2020. The SRC D water managers are available to provide assistance to landowners during the process of filing. If you have questions about filing your report, please contact your water manager.

Water Manager Office:
(707) 426-2431

Tim Edmunds - (707) 639-6689

Jeff Taylor - (707) 639-6690

Phelan McKinney - (707) 631-0819

tedmunds@suisunrcd.org

jtaylor@suisunrcd.org

pmckinney@suisunrcd.org

Preservation Implementation Act (PAI) cost share is available for projects upgrading corrugated metal pipes and cast iron gates to corrosive resistant material. The first round of PAI projects has been reviewed, and the deadline for submitting an application is June 19th.

SRC D is conducting interviews with managers at each club to update the Individual Management Plans last certified in the 1980s. As we continue to update the management plans, we will be contacting each club. Drafts are scheduled to be completed by the end of the summer.

Land of the West Wind

Quarterly newsletter of the Suisun Resource Conservation District
Address: 2544 Grizzly Island Road
Suisun, CA 94585
Telephone: (707) 425-9302
Fax: (707) 425-4402
Water Manager office: (707) 426-2431
E-mail: srcd@suisunrcd.org
Website: www.suisunrcd.org

SRCD Staff

Steven Chappell, *Executive Director*
John Takekawa, *Operations Manager*
Kelli Perez, *Office Supervisor*
Tim Edmunds, *Biologist/Water Manager*
Phelan McKinney *Biologist/Water Manager*
Jeff Taylor, *Biologist/Water Manager*
Jesirae Collins, *Biologist*
Mark Mouton, *Caretaker LJI*

Board of Directors

Tony Vaccarella, *President*
(650) 365-1642
Terry Connolly, *Finance*
(707) 4964-1105
Arnold Lenk, *Agency Relations*
(925) 284-3100
Jim Waters, *Legal*
(510) 409-3864
Mike Lewis, *Personnel*
(707) 224-3824

Associate Directors

Dennis Becker
Kurt Black
H. Kent Hansen
Steve Roerden
Lalo Kwiat

Directors Emeritus

(*deceased)
James Bancroft*
Paul Crapuchettes*
Ray Lewis*
Dr. William Coon*
Greg Palamountain*
Timothy Egan*
Leland Lehman*

SRCD's public meetings are held at 2 PM on the second Wednesday of each month at the Solano County Supervisors Chambers
675 Texas Street
Fairfield, CA 94533

SRCD represents private landowners of the Suisun Marsh at the Federal, State, and local levels. Its historic goal has been to achieve a water supply of adequate quality so that preferred wetland habitat values will be retained through appropriate management practices. Through cooperation with landowners and various agencies, SRCD seeks to develop new programs aimed at protecting and improving the Suisun Marsh for future generations.

Tracking Ducks to Your Duck Club

Mike Casazza, *Research Wildlife Biologist*
USGS Western Ecological Research Center

Starting in 2015, the USGS and the California Department of Water Resources (DWR) and SRCD began tracking common dabbling ducks using cell-phone tower (GSM) and GPS technology within Suisun Marsh. Suisun Marsh Program initiated a collaborative effort to better understand the value of Suisun Marsh for waterfowl within the Pacific Flyway. This effort is in support of DWR's 1984 Plan of Protection for the Suisun Marsh, the 2013 Suisun Marsh Habitat Management, Preservation, and Restoration Plan, and the 2015, the Suisun Marsh Preservation Agreement. These Plans and Agreements are intended to mitigate the effects of the Federal Central Valley Project and the State Water Projects on the Suisun Marsh and conserve its wetland and wildlife resource.



Hen with telemetry backpack

The tracking uses small solar powered and cell network connected GPS backpacks. Since 2015, nine duck species have been marked (Table 1). Though the marked birds make up a small fraction of the total birds using the Marsh, their location information is representative of the wider population and can provide useful insights to aid land and wildlife management. In an effort to make this information available to public and private land managers, USGS, DWR, and SRCD have developed waterfowl reports for each parcel in the Marsh.



Trapped ducks are banded and are selected for a transmitter

The goal of the reports is to provide land managers with information on the timing, precise locations, and species that are using the parcel of interest. This information could be used to help identify areas of successful management which could be replicated in other locations.

The reports include information describing the number of marked individuals and their locations in each parcel throughout the year. It is important to note that each location is likely representative of many birds. The year is broken into 4 time periods—Spring staging, Summer nesting and brooding (resident birds), Early hunting season, and Late hunting season—to help identify how habitat use changes over time. The early and late hunting season time periods are further broken into day and night use. The reports for each parcel will be available through the SRCD.

Tracking. Continued on page 3.

The Hill Slough Project Has Resumed


Sarah Estrella
Environmental Specialist,
California Department of Fish and Wildlife

Work has resumed along the northernmost mile of Grizzly Island Road. Raising this section is one of the first steps in the 850-acre Hill Slough Tidal Restoration Project at CDFW's Hill Slough Wildlife Area. Flooding during high tides and storms is an increasing problem with rising sea levels and raising this road is a required project component.

The first section of a 2-mile-long loop trail will be constructed next, at the northeast side, fol-



Hill Slough Project Plan



lowed by habitat improvements, and finally levee breaches by the end of 2021. This work will eliminate maintenance of 9 miles of levees, reduce mosquito habitat, and contribute to the recovery of species such as green sturgeon and Sacramento River winter-run Chinook salmon. The roadwork, a joint effort between CDFW and Solano County, continues by placing imported fill along the west and east sides of the road. Once the shoulders and side swales are filled to grade, the contractor will bring the roadway travel lanes up to final elevation in a series of lifts. Each lift will be leveled and traffic will run on dirt until this process is completed. Following this step, an aggregate base will cap the top. The contractor will then construct the travel lanes and allow the roadway section to settle through the summer, fall, and winter months. Travelers should anticipate minor delays and traveling on dirt surfaces. Reduced speed limits will be posted during this phase of work. Following the settlement period, when weather permits in 2021, the road will be completed and paved.



Alligator weed in Suisun Marsh: free-floating aquatic, and emergent floating-leaved mats

Dispersal: The plant spreads asexually with floating shoot fragments drifting with water currents. No seed production has been observed. The hollow stems break very easily, and even very small stem fragments can float to new sites and establish a new patch. Be a good neighbor: It is important to avoid stem breakage and prevent water dispersal of fragments that can easily spread the invasion.

Impacts: Alligator weed displaces open water habitat and desirable vegetation; degrades habitat and water quality; impedes water management, recreation, navigation; increases management costs.

Management: Aquatic-labelled glyphosate seems to be the most common herbicide used on alligator weed. Integrated weed management has been more effective than single herbicide management approaches: herbicide application followed by mechanical removal using floating booms to trap fragments (successful in Australia and elsewhere); biological control (insect herbivory) coupled with herbicide follow-up is extremely effective in Florida and throughout warm southern states at great cost savings over herbicides alone. USDA-ARS is seeking and evaluating insect agents tolerant of northern California climate. In NWR-managed seasonal wetlands, herbicide treatments of alligator weed and the response of desired native vegetation in the waterfowl habitat was assessed. Both Triclopyr amine and Imazapyr were effective at high rates of application in July; April applications were less effective. However, April applications of triclopyr were best for restoration of desired vegetation, while Imazapyr impeded succession and desirable vegetation (consistent with observations in California wetlands).

Current Treatment Programs: Currently, there is a moratorium on CA Boating and Waterway’s herbicide treatments for aquatic weeds in Suisun Bay and Suisun Marsh due to regulatory permits that limit target weeds, timing of applications and herbicides that can be used, and geographic scope. Permitted treatments of alligator weed are underway in the Delta using foliar spray applications of glyphosate with Agridex® surfactant administered from airboats or shallow draft motor boats. A pilot-scale field study was initiated in 2019 to evaluate efficacy of a Glyphosate-Imazamox-Competitor® surfactant mix (not currently permitted for operational use) as a potential improvement for control of alligator weed, and results will inform the biological assessment and permitting for resumed treatment of aquatic weeds in Suisun Bay and waterways of Suisun Marsh. This state management program is limited to treatments in public waterways, but future treatments can hopefully reduce dispersal of alligator weed into managed wetlands. Within wetland treatments should be coordinated with SRCD. Watch for alligator weed and inform SRCD of new infestations.


Don’t worry, no alligators have been spotted in Suisun Marsh, only alligator weed!

Telemetry Data From A Duck Club

How do the birds use the marsh? Telemetry data collected by the United States Geological Survey continues to help us answer this question. Using the small, light weight transmitters, researchers have been able to determine how far the ducks moved, how much space they used when not flying (e.g. when foraging or roosting), and how their time was allocated across the day. Results showed that the ducks may travel shorter distances and use smaller areas than previously thought, indicating that central valley ducks benefit from small, resource-rich habitat patches.



Species	2015	2016	2017	2018	2019	2020	Total
Blue-winged teal	0	0	0	4	0	0	4
Canvasback	0	8	3	0	0	0	11
Cinnamon teal	0	0	14	10	10	0	34
Eurasian wigeon	3	13	24	0	0	1	41
Gadwall	13	34	6	23	12	0	88
Green-winged teal	0	0	0	0	1	16	17
Mallard	34	65	38	27	24	1	189
Northern pintail	19	39	31	37	7	2	135
Northern shoveler	5	19	2	0	2	0	28

The Table above shows the number and species of marked birds that use Suisun Marsh.



Locations of marked birds on Lower Joice Island Duck Club.

Day and Night locations at Lower Joice Island in 2015 during early hunt season (September through November).



Lower Joice Island is characterized by differences in habitat between the north and the south half of the Island. The north features open water, often favored by pintail, while the southern ponds have more emergent vegetation, such as Alkali Bulrush. Telemetry data can help back assumptions that are widely accepted by duck hunters and biologists, or uncover new ideas. Telemetry data shows the special distribution broken down by species as what we might expect, but also has shown that birds move shorter distances during the season than previously thought, with most flights being less than a kilometer.


Juvenile Salmon Distribution, Abundance, and Growth in Restored and Relict Bay-Delta Marsh Habitats

Brett Harvey,
Senior Environmental Scientist
Department of Water Resources Division of Environmental Services

You may have wondered what the colorful floating squares were in Cross Slough, or noticed a trawling research vessel in Suisun Slough. These are part of a slough of exciting fish studies currently taking place in Suisun Marsh.

The study *Juvenile Salmon Distribution, Abundance, and Growth in Restored and Relict Bay-Delta Marsh Habitats* (aka Tidal Parr Study) is a group of collaborative studies underway in the Upper Estuary that began in January 2019, and will continue through June 2022. The studies are investigating how juvenile salmon use and benefit from marsh habitats of different qualities in the region from Sherman Lake to San Pablo Bay. Although this region is the location of much ongoing and planned habitat restoration, the way that juvenile salmon use these habitats is one of the least understood aspects of the salmon life cycle in the Central Valley. The information we collect will be used to better inform and guide restoration and resource management.

The study involves Trawl Surveys, Salmon eDNA Surveys, a Stable Isotope and Otolith Microstructure Study, and a Cage Growth Study. This year (2020), the Cage Growth Study included a side experiment: the manipulation of drain gates into Luco Slough to observe changes in water quality, aquatic bugs (fish food), and caged fish growth.




During the first year of the study, a “wet” year, we found that wild (non-hatchery) salmon were widely distributed across the study region, and that there were large differences in growth rates across cage locations. This year, a relatively “dry” year, was going to be an interesting comparison. Unfortunately the Covid-19 lockdown interfered with 2020 data collection for all studies. Useful data was still recovered and the comparison between wet and dry conditions will be of particular focus in future analyses. The study team hopes to be back in the field in full force for the final field season in 2021, and to repeat and expand the pond drain manipulation study.



Suisun Marsh Weed Alert: Alligator Weed!

Brenda J. Grewell, PhD
USDA-ARS Invasive Species and Pollinator Health Research Unit
Dept of Plant Sciences, UC Davis

Alligator weed (*Alternanthera philoxeroides*), a new uninvited weed, has arrived in the Suisun Marsh. This South American immigrant is considered one of the world’s most aggressive invasive plant species. It was introduced to the southeastern U.S. in about 1900, and aggressively spread throughout waterways and wetlands of southern states. Arrival in Los Angeles County was noted in 1940, and populations spread to Riverside, Tulare, Kings, and San Bernardino counties. California Dept. of Food and Agriculture designates alligator weed an A-rated Noxious Weed. A USDA-ARS research team reported the first northern CA observation of alligator weed in August 2017 in Montezuma Slough at Grizzly Island. The population in Suisun Marsh is genetically distinct from those in southern California, and therefore this is a new and recent introduction. State Park’s Division of Boating & Waterways quickly requested a risk assessment by California Dept. of Fish and Wildlife to facilitate permitting for management of the weed. Treatments were initiated in 2018, and new detections indicated a wider distribution in the Delta and Sacramento River.



California Department of Parks and Recreation
Division of Boating and Waterways



An Alligator Weed Workgroup convened and conducted blitz surveys in 2019. New detections documented spread in the Sacramento, San Joaquin and Feather Rivers, Delta and Suisun Marsh. The most impacted Suisun areas are Chipps Island, Van Sickle Island-Spoonbill Creek, Montezuma and Cutoff Sloughs.

Identification: Three best traits for i.d are: 1) opposite leaves, 2) hollow stem, 3) cluster of white flowers less than an inch in diameter on a short stalk at stem nodes or at terminal end of stem.



Flowers on short stalks



Small white flower cluster



Hollow stem and opposite leaves

Alligator Weed, Continued on page 6